The Suicide Behaviors Questionnaire-Revised (SBQ-R) - Overview

The SBQ-R has 4 items, each tapping a different dimension of suicidality:1

- Item 1 taps into lifetime suicide ideation and/or suicide attempt.
- Item 2 assesses the frequency of suicidal ideation over the past twelve months.
- Item 3 assesses the threat of suicide attempt.
- Item 4 evaluates self-reported likelihood of suicidal behavior in the future.

Clinical Utility

Due to the wording of the four SBQ-R items, a broad range of information is obtained in a very brief administration. Responses can be used to identify at-risk individuals and specific risk behaviors.

Scoring

See scoring guideline on following page.

Psychometric Properties¹

	Cutoff score	Sensitivity	Specificity
Adult General Population	≥7	93%	95%
Adult Psychiatric Inpatients	≥8	80%	91%

^{1.} Osman A, Bagge CL, Guitierrez PM, Konick LC, Kooper BA, Barrios FX., The Suicidal Behaviors Questionnaire-Revised (SBQ-R): Validation with clinical and nonclinical samples, Assessment, 2001, (5), 443-454.

SBQ-R - Scoring

Selected response 1	Non-Suicidal subgroup	1 point	
Selected response 2	Suicide Risk Ideation subgroup	2 points	
Selected response 3a or 3b	Suicide Plan subgroup	3 points	Section 1
Selected response 4a or 4b	Suicide Attempt subgroup	4 points	Total Points

Selected Response:	Never	1 point	
	Rarely (1 time)	2 points	
	Sometimes (2 times)	3 points	
	Often (3-4 times)	4 points	3
	Very Often (5 or more times)	5 points	Total Points

Item 3: taps into the threat of suicide attempt					
Selected response 1	1 point				
Selected response 2a or 2b					
Selected response 3a or 3b	3 points	Total Points			

Selected Response:	Never	0 points	
	No chance at all	1 point	
	Rather unlikely	2 points	
	Unlikely	3 points	
	Likely	4 points	
	Rather Likely	5 points	
	Very Likely	6 points	Total Points

Sum all the scores circled/checked by the respondents.

The total score should range from 3-18.

Total Score

AUC = Area Under the Receiver Operating Characteristic Curve; the area measures discrimination, that is, the ability of the test to correctly classify those with and without the risk. [.90-1.0 = Excellent; .80-.90 = Good; .70-.80 = Fair; .60-.70 = Poor]

	Sensitivity	Specificity	PPV	AUC
 Item 1: a cutoff score of ≥ 2 Validation Reference: Adult Inpatient Validation Reference: Undergraduate College 	0.80 1.00	0.97 1.00	.95 1.00	0.92 1.00
Total SBQ-R: a cutoff score of ≥7 • Validation Reference: Undergraduate College	0.93	0.95	0.70	0.96
Total SBQ-R: a cutoff score of ≥ 8 • Validation Reference: Adult Inpatient	0.80	0.91	0.87	0.89

SBQ-R Suicide Behaviors Questionnaire-Revised

Patient Name		ne			Date of Visit	
In	stru	ction	ns: Please check the number beside applies to you.	the s	tater	ment or phrase that best
1.	Hav	/e v	ou ever thought about or attemp	ted :	to ki	II yourself? (check one only)
	П	1.	Never			The state of the s
		2.	It was just a brief passing thought			
			I have had a plan at least once to k	ill mv	self l	but did not try to do it
			I have had a plan at least once to k	1950		\$250. 20 0000 As (Mail
	П		I have attempted to kill myself, but	10.50		R 001
			I have attempted to kill myself, and			
			, , , , , , , , , , , , , , , , , , , ,		,	
2.	Ηοι	∧ of	ten have you thought about killi	ng y	ours	elf in the past year? (check one only)
		1.	Never			
		2.	Rarely (1 time)			
		3.	Sometimes (2 times)			
		4.	Often (3-4 times)			
		5.	Very Often (5 or more times)			
3.	Hav	e yo	ou ever told someone that you w	ere g	going	g to commit suicide,
	or t	hat	you might do it? (check one only)			
		1.	No			
		2a.	Yes, at one time, but did not really	want	to d	ie
		2b.	Yes, at one time, and really wanted	to di	e	
		За.	Yes, more than once, but did not w	ant t	o do	it
		3b.	Yes, more than once, and really war	nted	to do	o it
4.	Hov	low likely is it that you will attempt suicide someday? (check one only)				
		0.	Never		4.	Likely
		1.	No chance at all		5.	Rather likely
		2.	Rather unlikely		6.	Very likely
		3.	Unlikely			